



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

the city bacteriologist for examination, after being dipped in petrol to kill the fleas. Since the beginning of the current year more than 5,500 rats have been caught, of which 3,000 were examined and of this number only three were found plague infected. One of these was caught on board ship April 30 and the other two were caught in adjacent warehouses August 1 and 30, respectively.

A rat catcher is put on board every ship coming from a port suspected of being infected with plague and all precautions are taken to prevent rats coming ashore. The area in which the two infected rats were found is about 2 miles from the transatlantic liners' docks. During the past eight or nine years about 35,000 rats have been caught in the port area and forwarded to the university for examination and in no case was plague discovered.

HAWAII.

Examination of Rodents for Plague Infection.

During the week ended September 14, 1912, 593 rats and mongoose were examined at Hilo and 1,596 at Honokaa. No plague infection was found.

At Honolulu during the same week 166 rats were examined. No plague infection was found.

The last case of human plague occurred at Pepeekeo, Hawaii, September 13, 1912. The last plague-infected rat was found at Olaa, Hawaii, September 9, 1912.

ITALY.

Genoa—Epidemic Typhoid Fever.

An epidemic of typhoid fever has been reported present in Genoa. From August 31 to September 26, 1912, there were reported 704 cases with 46 deaths.

JAPAN.

Cholera.

Surg. Irwin at Yokohama reports September 16 and 23: Cholera is spreading in southern Japan. At Osaka 12 cases were reported from September 14 to 21; at Moji 142 cases to September 21. At Shimonoseki, opposite Moji, 41 cases have been reported. On September 19, 4 cases were reported at Kyoto. On September 20 a total of 231 cases of which 136 were in Fukuoka-ken and 57 in Yamaguchi-ken was reported for all Japan. On September 23 the number of cases reported present in Fukuoka-ken was stated to be 330.

Two cases of cholera with one death were reported September 12 and 18 respectively, at Kobe.

Epidemic Cerebrospinal Meningitis.

Cerebrospinal meningitis is reported present in epidemic form at Okayama. On September 19 there were reported 21 new cases, making 347 cases from the date of the outbreak.